

Willow Pediatrics emphasizes the importance of preventative care, early identification of abnormalities in growth or development, and relationships with families that are built on mutual trust. For these reasons, we strongly encourage families to schedule well child visits as outlined below and to view the applicable CDC vaccine information sheets and vaccine package inserts on our [resource page](#). There are also age-based developmental handouts available. Please review the documents that are relevant to your child.

We also have specific intake forms for each well-child visit. Please have any requested forms completed prior to your visit. These forms allow us to prepare any additional material that may be helpful to your family, as well as tailor the appointment time to the unique needs of your family.

Age	Immunizations	Labs	Screenings
Newborn	Hep B #1		
2 weeks			
1 month	Hep B #2		Edinburgh Postnatal Depression Scale (EPDS) Perinatal Anxiety Screening Scale (PASS)
2 months	Pentacel #1 (DTaP/Polio/Hib) Pprevnar #1 RotaTeq #1 (given orally)		Edinburgh Postnatal Depression Scale (EPDS) Perinatal Anxiety Screening Scale (PASS)
4 months	Pentacel #2 (DTaP/Polio/Hib) Pprevnar #2 RotaTeq #2 (given orally)		Edinburgh Postnatal Depression Scale (EPDS) Perinatal Anxiety Screening Scale (PASS)
6 months	Pentacel #3 (DTaP/Polio/Hib) Pprevnar #3 RotaTeq #3 (given orally) Flu vaccine #1 *		Edinburgh Postnatal Depression Scale (EPDS) Perinatal Anxiety Screening Scale (PASS)
9 months	Hep B #3 Flu vaccine #2 *	Hemoglobin Lead	Ages and Stages Questionnaire (ASQ)
12 months	MMR #1 Hib #4		
15 months	Varivax #1 (chicken pox) Hep A #1	Hemoglobin Lead	
18 months	DTaP #4 Pprevnar #4		Ages and Stages Questionnaire (ASQ) M-CHAT

2 years	Hep A #2	Hemoglobin Lead	M-CHAT
2 1/2 years			
3 years		Hemoglobin Lead	Vision Hearing
4 years	MMR #2		Vision Hearing
5 years	Varivax #2 (chicken pox)		Vision Hearing
6 years	Polio #4 DTaP #5		Vision Hearing
7 years			
8 years			Vision Hearing
9 years			
10 years		Lipids	Vision Hearing Scoliosis (females)
11 years	Menactra #1 Tdap #1 Gardasil #1		
12 years	Gardasil #2		Vision Hearing Depression Scoliosis (females)
13 years			Depression
14 years			Depression Scoliosis (males)
15 years			Vision Hearing Depression
16 years	Menactra #		Depression
17 years		Lipids HIV	Depression
18 years			Hearing

			Depression
19 years			Depression
20 years			Depression
21 years			Depression

*** Please note:** The influenza vaccine (flu vaccine) is given annually, starting at 6 months of age. In order to be optimally protected, the first year your child receives the flu vaccine he or she will need 2 doses given 1 month apart. Thereafter, your child will need one dose yearly. Flu vaccines are usually available annually between August and February each year, while supplies last.

Client Signature

Date